

MEDICATION DEFERRAL LIST

DO NOT STOP taking medications prescribed by your doctor in order to donate blood. Donating while taking these drugs could have a negative effect on your health or the health of the recipient of your blood.

(REFER TO QUESTION #4 ON THE DONATION RECORD)

SOME MEDICATIONS MAY AFFECT YOUR ELIGIBILITY TO DONATE BLOOD.

PLEASE TELL US IF YOU...

Are being treated with the following types of medications...	or have taken...		which is also called...	anytime in the last...
Anti-platelet agents (usually taken to prevent stroke or heart attack)	Feldene		piroxicam	2 days
	Effient		prasugrel	3 days
	Brilinta		ticagrelor	7 days
	Plavix		clopidogrel	14 days
	Ticlid		ticlopidine	
	Zontivity		vorapaxar	1 month
Anticoagulants or “blood thinners” (usually to prevent blood clots in the legs and lungs and to prevent strokes)	Arixtra		fondaparinux	2 days
	Eliquis		apixaban	
	Fragmin		dalteparin	
	Pradaxa		dabigatran	
	Savaysa		edoxaban	
	Xarelto		rivaroxaban	
	Coumadin Warfilone Jantoven		warfarin	7 days
	Lovenox		enoxaparin	
	Heparin, low-molecular-weight heparin			
Acne treatment	Accutane Amnesteem Absorica Claravis Myorisan Sotret Zenatane		isotretinoin	1 month
Multiple myeloma	Thalomid		thalidomide	1 month
	Revlimid		lenalidomide	
Rheumatoid arthritis	Rinvoq		upadacitinib	1 month
Hair loss remedy	Propecia		finasteride	1 month
Prostate symptoms	Proscar		finasteride	1 month
	Avodart or Jalyn		dutasteride	6 months
Immunosuppressant	Cellcept		Mycophenolate mofetil	6 weeks
	Myfortic		Mycophenolic Acid	
Hepatitis exposure	Hepatitis B Immune Globulin		HBIG	3 months
HIV Prevention (also known as PrEP and PEP)	Any medication taken by mouth (oral) to prevent HIV.	Truvada	emtricitabine and tenofovir disoproxil fumarate	3 months
		Descovy	emtricitabine and tenofovir alafenamide	3 months
		Injectable HIV Prevention	Apretude	cabotegravir
Basal cell skin cancer	Erivedge		vismodegib	2 years
	Odomzo		sonidegib	
Relapsing multiple sclerosis	Aubagio		teriflunomide	2 years
Rheumatoid arthritis	Arava		leflunomide	2 years
Psoriasis	Soriatane		acitretin	3 years
	Tegison		etretinate	Ever
HIV Treatment	Any medication to treat HIV. May also be called antiretroviral therapy (ART)			Ever
Experimental Medication or Unlicensed (Experimental) Vaccine				12 months

MEDICATION DEFERRAL LIST

Some medications may affect donor eligibility for the following reasons:

Antiplatelet agents affect platelet function, so people taking these drugs should not donate platelets for the indicated time. You may still be able to donate whole blood or red blood cells by apheresis.

Anticoagulants or "blood thinners" are used to treat or prevent blood clots in the legs, lungs, or other parts of the body, and to prevent strokes. These medications affect the blood's ability to clot, which might cause excessive bruising or bleeding when you donate. You may still be able to donate whole blood or red blood cells by apheresis.

Isotretinoin, finasteride, dutasteride, acitretin and etretinate can cause birth defects. Your donated blood could contain high enough levels to damage the unborn baby if transfused to a pregnant woman.

Thalomid (thalidomide), Revlimid (lenalidomide), Erivedge (vismodegib), Odomzo (sonidegib), Aubagio (teriflunomide) and Rinvoq (upadacitinib) may cause birth defects or the death of an unborn baby if transfused to a pregnant woman.

Cellcept (mycophenolate mofetil) and Arava (leflunomide) are immunosuppressants that may cause birth defects or the death of an unborn baby if transfused to a pregnant woman.

PrEP or pre-exposure prophylaxis involves taking a specific combination of oral medicines (i.e., short-acting antiviral PrEP) or injections (i.e., long-acting antiviral PrEP) as a prevention method for people who are HIV negative and at high risk of HIV infection. FDA has determined that the available data demonstrate that the use of PrEP or PEP may delay the detection of HIV by currently licensed screening tests for blood donations, potentially resulting in false negative results in infected individuals. Although "Undetectable = Untransmittable" for sexual transmission, this **does not apply to transfusion transmission**.

PEP or post-exposure prophylaxis is a short-acting treatment started as soon as possible after a high-risk exposure to HIV to reduce the risk of infection. FDA has determined that the available data demonstrate that the use of PrEP or PEP may delay the detection of HIV by currently licensed screening tests for blood donations, potentially resulting in false negative results in infected individuals. Although "Undetectable = Untransmittable" for sexual transmission, this **does not apply to transfusion transmission**.

ART or antiretroviral therapy is the use of a combination of HIV medicines (called an HIV regimen) to treat HIV infection. HIV infection requires a permanent deferral despite treatment with ART. Antiretroviral drugs do not fully eliminate the virus from the body and donated blood from individuals infected with HIV taking ART can potentially still transmit HIV to a transfusion recipient. Although "Undetectable = Untransmittable" for sexual transmission, this **does not apply to transfusion transmission**.

Hepatitis B Immune Globulin (HBIG) is an injected material used to prevent hepatitis B infection following a possible or known exposure to hepatitis B. HBIG does not prevent hepatitis B infection in every case; therefore, persons who have received HBIG must wait to donate blood.

Experimental Medications are usually associated with a research study and the effect on the safety of transfused blood is unknown.